

Universalities in One-electron Properties of Limit Quasi-periodic Lattices

Rihei Endou^{1,*}, Komajiro Niizeki¹, and Nobuhisa Fujita²

¹*Department of Physics, Graduate School of Science, Tohoku University, Sendai 980-8578, Japan and*

²*Structural Chemistry, Arrhenius Laboratory, Stockholm University, 10691 Stockholm, Sweden*

(Dated: February 2, 2008)

We investigate one-electron properties of one-dimensional self-similar structures called limit quasi-periodic lattices. The trace map of such a lattice is nonconservative in contrast to the quasi-periodic case, and we can determine the structure of its attractor. It allows us to obtain the three new features of the present system: 1) The multi-fractal characters of the energy spectra are *universal*. 2) The supports of the $f(\alpha)$ -spectra extend over the whole unit interval, $[0, 1]$. 3) There exist marginal critical states.

PACS numbers: 61.44.Br, 64.60.Ak, 71.23.Ft

Since the discovery of a quasicrystal, deterministic and aperiodic systems have been attracted much attention [1]. They are classified into the third group in addition to periodic systems and random systems. Although they include a wide range of diverse structures, a self-similar structure such as the Fibonacci lattice or the Penrose tiling is especially important because quasicrystals belong to materials of this type. In particular, a one-dimensional (1D) self-similar structure has the symmetry described by a semi-group, and the electronic state on it exhibits rich properties [2, 3, 4, 5]. The presence of multifractal energy spectra and self-similar wave functions is peculiar to such systems. Here we will add *universality* to the subject; it applies to limit quasiperiodic lattices which form an important class of self-similar structures.

A 1D self-similar structure is produced with a substitution rule (SR) starting from two elements, A and B [6]. It is taken to be an aperiodic lattice composed of two types of sites, and called a self-similar lattice (SSL). A general form of the SR is written as $A \rightarrow \sigma_A(A, B)$, $B \rightarrow \sigma_B(A, B)$, where $\sigma_A(A, B)$ and $\sigma_B(A, B)$ are words of the two letters or, equivalently, finite sequences of the two symbols. The SR can be specified by a combination, $(\sigma_A(A, B), \sigma_B(A, B))$, of two different words, and an infinite number of different SRs are formed by changing the combination. A simplest SSL is the Fibonacci lattice, whose SR is (B, AB) , while another one with the SR, $(B, BAAB)$, produces the mixed mean lattice (MM lattice) [7]. To every SR, (σ_A, σ_B) , we can associate a Frobenius matrix, $M := \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ with a and c (resp. b and d) being the numbers of A and B in $\sigma_A(A, B)$ (resp. $\sigma_B(A, B)$); we shall call it the substitution matrix of the SR or of the relevant SSL. An SR is not uniquely determined by its substitution matrix because two or more choices are usually possible for the orders of the letters, A and B , in σ_A and σ_B . If there are several locally-isomorphic classes of SSLs with a common substitution matrix, we may call them iso-

mers. For example, two SSLs with SRs, $(B, ABBA)$ and $(B, ABAB)$, are isomers of the MM lattice. When an SSL has a global center of the reflection symmetry, it can be called a symmetric SSL and the relevant SR a symmetric SR [8]. We may call an SR, (σ_A, σ_B) , palindromic if both σ_A and σ_B are palindromes, while we may call it quasi-palindromic if it is written with a palindromic SR, (σ'_A, σ'_B) , as $(\sigma'_A B, \sigma'_B B)$. Palindromic SRs and quasi-palindromic ones produce SSLs. If a substitution matrix is specified, at least one of the relevant isomers is a symmetric SSL. All the SRs presented above produce symmetric SSLs.

We shall turn to the structure factor $S(Q)$ of an SSL [9, 10]. The set of all the SSLs is divided into two classes Π and Γ ; $S(Q)$ for the case Π consists only of Bragg peaks, while the one for the case Γ is singular continuous. The class Π is divided, further, into the three subclasses, i) Π_I : Quasiperiodic, ii) Π_{II} : Limit Quasiperiodic, and iii) Π_{III} : Limit Periodic. Representatives of the three are the Fibonacci lattice, the MM lattice, and the period-doubling lattice, respectively. Let M be the substitution matrix of an SSL, $\mathcal{L}$, and $\tau (> 1)$ be the Frobenius eigenvalue of M . Then, a necessary and sufficient condition for $\mathcal{L}$ to belong to Π_I or Π_{II} is that the second eigenvalue τ' satisfies $|\tau'| < 1$. If it is satisfied, τ is a quadratic irrational called a Pisot number. Note that τ' is the algebraic conjugate of τ and $\tau\tau' = s$ with $s := \det M$; M is unimodular only when $s = \pm 1$. It is known that $\mathcal{L}$ belongs to Π_I if $|s| = 1$ but to Π_{II} otherwise. For example, $\tau = (1 + \sqrt{5})/2$ (resp. $\tau = 1 + \sqrt{3}$) for the Fibonacci lattice (resp. the MM lattice). The set of all the wave numbers at the peaks of $S(Q)$ form an additive group, $\mathcal{M}$, which is called the Fourier module. The two subclasses Π_I and Π_{II} are distinguished by whether the number of the generators of $\mathcal{M}$ is two or infinite. Let M be the substitution matrix of an SSL and let $\omega := (\tau - a)/b = c/(\tau - d)$ with τ being the Frobenius eigenvalue of M . Then, the Frobenius eigenvector of M is given by the column vector ${}^t(1, \omega)$. The Fourier module $\mathcal{M}$ of an SSL in Π_{II} is written as $\mathcal{M} = (1 + \omega)^{-1} \mathbf{Z}\{\omega\}$ with $\mathbf{Z}\{\omega\} := \mathbf{Z}[\omega] \cup \tau^{-1} \mathbf{Z}[\omega] \cup \tau^{-2} \mathbf{Z}[\omega] \cup \dots$, where $\mathbf{Z}[\omega] := \{x + y\omega \mid x, y \in \mathbf{Z}\}$. The Fourier module is a dense set on the real axis. Note that $\mathcal{M}$ is common among dif-

*Electronic address: rihei@cmpt.phys.tohoku.ac.jp




FIG. 1: The energy spectrum of the MM lattice with $\Delta := V_B - V_A = 1$

ferent isomers because $\mathcal{M}$ is uniquely determined by M . Incidentally, $\mathbf{Z}\{\omega\} = \mathbf{Z}[\omega]$ and $\mathcal{M} = (1 + \omega)^{-1}\mathbf{Z}[\omega]$ for an SSL in Π_I because $\tau\mathbf{Z}[\omega] = \mathbf{Z}[\omega]$ for this case.

Every set, say Π , of SSLs is divided into the symmetric subset Π^s and its complement Π^a , where Π^s is composed only of symmetric SSLs. Previous researches on the one-electron states of SSLs are focused exclusively on the case of class Π^s . More precisely, the subclass Π_I^s is investigated extensively [5] but Π_{II}^s is investigated by nobody, while there are a considerable number of papers on the subclass Π_{III}^s but they are still on a rudimentary stage (see, for example, [7]). Therefore, the authors investigated in detail the case of the subclass Π_{II}^s . Though Π_{II}^s appears a simple extension of Π_I^s , it turns out as shown below that its one-electron states exhibit a remarkably different character from the case of the latter. We shall call hereafter an SSL in Π_{II}^s a limit quasi-periodic lattice (LQL).

Self-similarity of an SSL allows us to use the trace map (TM) for researches of the electronic states on it [2, 3]. The character of the TM as a non-linear map depends crucially on whether it is conservative or not [7, 11, 12]. It is known that SSLs with conservative TMs form a special subset, Π_I^c , of Π_I^s [8, 12]. Hereafter, an SSL in Π_I^c is called simply a conservative SSL (CSSL). Remember that an LQL is never conservative.

Generic characters of one-electron states of an SSL can be investigated using the tight-binding Hamiltonian $\mathcal{H}$, each site-energy of which takes V_A or V_B depending on the type of the site [5]. The unit of energy can be so chosen that the transfer integral assumes -1 . It was shown generally that the energy spectrum $\sigma = \sigma(\mathcal{H})$ is singular continuous [13]. More precisely, it is a bounded closed set having an infinite number of gaps, and, moreover, the length of its every connected component vanishes; we can see it in Fig.1. This means that the integrated state density $H(E)$ per one site is a devil's staircase. Every gap of σ can be specified by the height of the corresponding step of the staircase. Let G_σ be the set formed by adding 0 and 1 to the set of the heights of all the steps. Then, it is a discrete subset of the unit interval $[0, 1]$, and is determined by the gap labeling theorem [14]. Since σ is a multifractal, its singularity, $\alpha = \alpha(E)$ with $E \in \sigma$, is a function of E . The distribution of α in σ is characterized by the $f(\alpha)$ -spectrum [5]. The support of $f(\alpha)$ is a closed interval, $[\alpha_{\min}, \alpha_{\max}] \subset [0, 1]$. For every E in σ , we can define a function by $h(\varepsilon) := H(E + \varepsilon/2) - H(E - \varepsilon/2)$. If

it satisfies the equation, $h(\varepsilon) \approx \mu h(\varepsilon/\lambda)$, for two parameters $\lambda = \lambda(E)$ and $\mu = \mu(E)$ chosen suitably, E is a center of local self-similarity of σ . The singularity α of the center is given by $\alpha = \ln \mu / \ln \lambda$ [3, 5]. A remarkable feature of an SSL is that the set, σ_{ls} , of all such centers is a countably infinite set [4]. The set, $\Sigma := \{H(E) | E \in \sigma_{ls}\}$ is an important object which characterizes σ_{ls} .

In the trace map formalism, a homomorphism is defined from $\mathcal{F}(A, B)$, the free semi-group generated by the two generators A and B , into $SL(2, \mathbf{R})$, a group formed of 2×2 unimodular matrices [7, 11, 16]. If $\hat{A}$ and $\hat{B}$ are the images of A and B in the homomorphism, the image, $W(\hat{A}, \hat{B})$, of a sequence $W(A, B)$ is called the transfer matrix associated with the sequence. We can form an infinite series of pairs, $\{A_n, B_n\}$, of sequences by using the recursion relation $\{A_{n+1}, B_{n+1}\} = \{\sigma_A(A_n, B_n), \sigma_B(A_n, B_n)\}$ supplemented by the initial condition, $\{A_1, B_1\} := \{A, B\}$. The corresponding pair, $\{\hat{A}_n, \hat{B}_n\}$, of transfer matrices satisfy a homologous recursion relation. Let $\{\hat{A}, \hat{B}\}$ be such a pair at any generation and $W(A, B)$ any sequence. Then, $\text{Tr } W(\hat{A}, \hat{B})$ is a polynomial with integral coefficients of the three variables, $x := \text{Tr } \hat{A}$, $y := \text{Tr } \hat{B}$, and $z := \text{Tr } (\hat{A}\hat{B})$. Thus, a map is defined from $\mathcal{F}(A, B)$ into $\mathbf{Z}[x, y, z]$. Let T_x , T_y , and T_z be the three polynomials defined as the images of the three sequences, $\sigma_A(A, B)$, $\sigma_B(A, B)$, and $\sigma_A(A, B)\sigma_B(A, B)$, respectively. Then, the trace map T which yields a 3D dynamical system over a 3D phase space $\mathbf{R}^3$ is defined by $\mathbf{r} \rightarrow T\mathbf{r} := (T_x(\mathbf{r}), T_y(\mathbf{r}), T_z(\mathbf{r}))$ with $\mathbf{r} = (x, y, z)$ [2, 6, 11]. For example, $T\mathbf{r}$ assumes $(y, x, yz - x)$ for the Fibonacci lattice but

$$(y, xyz - x^2 - y^2 + 2, x(y^2 - 1)z - x^2y - (y^2 - 3)y) \quad (1)$$

for the MM lattice. Noncommutativity between the transfer matrices in a pair $\{\hat{A}, \hat{B}\}$ can be quantified by the polynomial defined by [2]

$$I := \frac{1}{2}\text{Tr}[\hat{A}, \hat{B}]^2 = x^2 + y^2 + z^2 - xyz - 4, \quad (2)$$

which vanishes if they are commutative. It can be shown readily that $I(\mathbf{r}_1) = (\Delta/2)^2$. If the subscript n being suppressed at the moment is recovered, we may write $\mathbf{r}_{n+1} = T\mathbf{r}_n$. An orbit $O(E) := \{\mathbf{r}_n | n \in \mathbf{N}\}$ in the phase space is generated from the initial point $\mathbf{r}_1 = (V_A - E, V_B - E, (V_A - E)(V_B - E) - 2)$. An important property of the TM is that $T\mathbf{r}$ itself does not depend on the parameters V_A , V_B , and E .

It can be shown generally that there exists a polynomial $P(\mathbf{r})$ such that $[\hat{A}_{n+1}, \hat{B}_{n+1}] = P(\mathbf{r}_n)[\hat{A}_n, \hat{B}_n]$ for a palindromic SR but $[\hat{A}_{n+1}, \hat{B}_{n+1}] = P(\mathbf{r}_n)[\hat{A}_n, \hat{B}_n]\hat{B}_n$ for a quasi-palindromic one [17]. Moreover, $|P(\mathbf{r})| \equiv 1$ for the conservative case but $\deg(P) \geq 1$ otherwise. We obtain $P = z$ for the MM lattice, for example. It yields the identity $I(T\mathbf{r}) \equiv [P(\mathbf{r})]^2 I(\mathbf{r})$. Therefore, $I(\mathbf{r})$ is an invariant of the TM only for the case of a CSSL but $I(\mathbf{r})$ is a semi-invariant for the general case because the TM never changes its sign [6].

Let us assume that $\hat{A}_n$ and $\hat{B}_n$ for $\exists n \in \mathbf{N}$ are not commutative and, besides, that $P(x_n, y_n, z_n) = 0$ for some $E \in \sigma$. Then, $\hat{A}_m, \hat{B}_m$ are commutative for $\forall m \geq n+1$, and we can conclude that the relevant eigenstate will be extended as in the case of a periodic system [18]. Different extended states may be obtained from the roots of a similar equation at a different generation. Therefore, an SSL may have an infinite number of extended states unless it is not an CSSL.

The dynamical system defined by the TM, T , is characterized by its limit cycles (LCs). If a point in $\mathbf{R}^3$ belongs to an LC with period p , it is a fixed point of T^p and an orbit starting from it is a pure cycle. A necessary and sufficient (N&S) condition for $E \in \sigma$ to belong to σ_{ls} is that $O(E)$ falls on an LC. If this is satisfied, we can determine the scaling parameter λ by a linear analysis of T^p around the relevant fixed point, where p is the period of the LC, so that the singularity $\alpha = \alpha(E)$ can be evaluated [3]. The wave function of the corresponding eigenstate is known to be asymptotically self-similar; the spatial ratio of the self-similarity is equal to τ^p [5].

Though $I(x, y, z)$ is not an invariant of the TM, T , of an LQL, the curved surface defined by the equation $I(x, y, z) = 0$ is an invariant surface. Its restriction, $\mathbf{S}$, into the cube $[-2, 2]^3 \in \mathbf{R}^3$ is a closed surface, which is a ball with the tetrahedral point symmetry but has four cusps at $(2, 2, 2)$, $(2, -2, -2)$, $(-2, 2, -2)$, and $(-2, -2, 2)$ (see Fig.1c in [11]). A coordinate system can be introduced into $\mathbf{S}$ by $x = 2 \cos(2\pi u)$, $y = 2 \cos(2\pi v)$, and $z = 2 \cos[2\pi(u+v)]$ with (u, v) being a variable on the 2D torus $\mathbf{T}^2 := \mathbf{R}^2/\mathbf{Z}^2$. Actually, $\mathbf{S}$ is doubly covered by $\mathbf{T}^2$ because $\pm(u, v)$ are mapped onto a single point on $\mathbf{S}$. The origin $(0, 0)$ of $\mathbf{T}^2$ corresponds to $(2, 2, 2)$ in $\mathbf{R}^3$. The TM induces a 2D dynamical system (DS) over $\mathbf{T}^2$: $T(u, v) = (u, v)M$ with M being the relevant substitution matrix. With these machineries together with a theory of algebraic number theory, we can determine all the cycles of the 2D DS, which will be presented elsewhere.

For a periodic system ($\Delta = 0$), every 3D orbit of T is confined to $\mathbf{S}$ from the very beginning. The initial state $\mathbf{r}_1$ of the orbit $O(E)$ with $E = -2 \cos 2\pi\kappa$ corresponds to $(u_1, v_1) = \pm(\kappa, \kappa)$ on $\mathbf{S}$, where κ is the rationalized wave number of a plane wave state. It follows that $(u_n, v_n) = \pm(L_n^{(A)}\kappa, L_n^{(B)}\kappa)$ with $L_n^{(A)}$ and $L_n^{(B)}$ being the numbers of the letters in A_n and B_n , respectively. A N&S condition for the 2D orbit $\{(u_n, v_n) | n \in \mathbf{N}\}$ to converge on $(0, 0) \in \mathbf{S}$ is equivalent to $\kappa \in \mathcal{M}$ [9, 10]. On the other hand, the 2D orbit falls on a limit cycle with period p if and only if $\{(u_{n+p} \mp u_n, v_{n+p} \mp v_n) | n \in \mathbf{N}\}$ converges on $(0, 0) \in \mathbf{S}$. Using the fact that the ratios $L_{n+p}^{(A)}/L_n^{(A)}$ and $L_{n+p}^{(B)}/L_n^{(B)}$ tend to τ^p as n goes to the infinity, we can show that a N&S condition for that condition to be satisfied is that $(\tau^p - 1)\kappa \in \mathcal{M}$ or $(\tau^p + 1)\kappa \in \mathcal{M}$. It follows that a N&S condition for the 2D orbit to fall on a limit cycle is given by $\kappa \in \mathbf{Q}[\omega] - \mathcal{M}$ with $\mathbf{Q}[\omega] := \{u + v\omega | u, v \in \mathbf{Q}\}$. Thus, the behavior of the TM of a periodic system has been completely

revealed. Since the energy spectrum of the periodic system is absolutely continuous, $\alpha(E)$ takes 1 [15]. Two states with wave numbers $\pm\kappa$ degenerate in energy and the corresponding two orbits of the TM are identical, so that we can assume $0 \leq \kappa \leq 1/2$. The number κ is nothing but the normalized value per one site of the number of the nodes in the relevant sinusoidal wave. Since $H(E)$ is related to $\kappa = \kappa(E)$ by the equation $H = 2\kappa(E)$, the asymptotic behavior of the TM at the energy E is determined by the value of $H(E)$, which belongs to the interval $[0, 1]$.

We will proceed to the case of a non-periodic system ($\Delta \neq 0$). Since an LQL has the reflection symmetry, every energy level has its own parity with respect to the center of symmetry. If Δ is changed adiabatically from zero to the present value, every energy level will change continuously because the degeneracies being present at $\Delta = 0$ does not matter if the parity is specified. It follows that κ remains to be the normalized number of the nodes. Then, the asymptotic behavior of the TM will be determined by the value $H = H(E)$, and the rule is the same as what is described above for the periodic system. Thus, we can conclude that $\Sigma = \mathbf{Q}[\omega] \cap [0, 1]$, which is a similar result to the one obtained in [4] for the case of the Fibonacci lattice. A remarkable property of the TM of an LQL is its *universality*. It is based on the two points: 1) The TM does not explicitly depend on the parameters Δ and E . 2) It is essentially of 3D character, and not confined to a 2D manifold as in the case of a CSSL. Thus, the attractor is independent of the value of Δ , and identical to that of the periodic system. It follows that the $f(\alpha)$ -spectrum of σ is universal. Moreover, we can conclude that $\alpha = 1$ for every center of local self-similarity of σ [15]. For example, a 3D orbit for the MM lattice converges to a 1-cycle if $\kappa = 1/3$, and the scaling parameter λ determined by a linear analysis of T around the relevant fixed point, $(-1, -1, -1)$, is found exactly equal to τ , so that $\alpha = 1$. As a consequence, we obtain $\alpha_{\max} = 1$.

From a brief perturbative consideration, we may expect that $G_\sigma = \mathcal{M} \cap [0, 1]$, which is guaranteed by the gap-labeling theorem [14]. The gap labeled by $H \in G_\sigma$ is bounded by two energy levels located at band edges. However, the corresponding fixed point $(2, 2, 2)$ of the TM is a singularity of $\mathbf{S}$, and we cannot conclude that $\alpha = 1/2$ for these two energy levels. Surprisingly, these states are actually marginal critical states, which we can prove by a similar technique to the one used in [19]. It follows that $\alpha_{\min} = 0$. This and the result $\alpha_{\max} = 1$ are consistent with that the $f(\alpha)$ -spectrum is universal because there is no reason why the value of $\alpha_{\min}$ or $\alpha_{\max}$ must take an odd value. The broadness of the support of the $f(\alpha)$ -spectrum means that the energy spectrum of an LQL is more inhomogeneous than that of a CSSL, which one can feel if he compares Fig.1 with the energy spectrum of, say, the Fibonacci lattice [5]. We show in Fig.2 the $f(\alpha)$ -spectrum of the energy spectrum in Fig.1.

If there are several isomers, their TMs are different.




FIG. 2: The $f(\alpha)$ -spectra of the energy spectra of periodic approximants for the MM lattice. The data from the 4-th to the 9-th generations are used. The data in the region $\alpha < .2$ are cut because their accuracies are insufficient on account of the presence of marginal critical states. The inset is a log-log plot of $\delta_n := \alpha_{\max}^{(n)} - 1$ versus the generation number n . The asymptotic linearity of the plot concludes that δ_n tends to zero in an inverse power law for n as n goes to the infinity. The $f(\alpha)$ -spectra of the Fibonacci lattice is superposed for a comparison.

Their $f(\alpha)$ -spectra belong, however, to a common universality class because each universality class of the $f(\alpha)$ -

spectrum of the LQL can be specified by the substitution matrix.

The TM derived from a CSSL is a 2D map on the curved surface $I(x, y, z) = (\Delta/2)^2$. Since this surface depends on the value of Δ , the $f(\alpha)$ -spectrum as well as $\alpha_{\min}$ and $\alpha_{\max}$ are not universal. This is consistent with the known fact that $\alpha_{\min} > 0$ and $\alpha_{\max} < 1$ for every CSSL (see Fig.2) [2]. The SSLs in the set Π_1^c are actually minorities in Π_1^s , and one-electron properties of SSLs in $\Pi_1^s - \Pi_1^c$ has never been investigated yet. The present theory is basically applicable to this case; the only modification needed is to replace the Fourier module by $\mathcal{M} = (1 + \omega)^{-1} \mathbf{Z}[\omega]$. Thus, among isomers belonging to Π_1^s , only the CSSL member is nonuniversal. Incidentally, the SR, (ABA, ABABA), produces a CSSL but its isomer, (ABA, BAAAB), does not.

Although no substance which takes a structure represented by an LQL as its thermodynamically stable phase is known, such a structure is realizable as an artificial super-lattice. Since a transfer matrix can be used also to a 1D Schrödinger equation, the results of this letter apply to the electronic state in such an artificial super-lattice as well[20]. Furthermore, it is basically applicable to a propagation in a stratified substance of an ultrasonic wave, an electromagnetic wave (light), and a spin wave as well. If, for example, a lower marginal critical state is used as a channel of light, the speed of light will fall dramatically.

In conclusion, one-electron properties of limit quasi-periodic lattices exhibits a universality in contrast to the case of quasi-periodic ones, and a universality class is specified solely by the substitution matrix.

One of the authors (K. N.) is grateful to Y. Morita.

-
- [1] *Physical Properties of Quasicrystals*, edited by Z. M. Stadnik, Series in Solid-State Sciences, 126 (Springer-Verlag, 1999).
 - [2] M. Kohmoto, P. Kadanoff, and C. Tang, Phys. Rev. Lett. **50**, 1870 (1983).
 - [3] M. Kohmoto and Y. Oono, Phys. Lett. **102A**, 145 (1984).
 - [4] P. A. Kalugin, A. Yu Kitaev, and L. C. Levitov, Pis'ma Zh. Eksp. Teor. Fiz. 41, 119 (1985) [JETP Lett. 41, 119 (1985)].
 - [5] For a review, see H. Hiramoto and M. Kohmoto, Int J. Mod. Phys. B**6**, 281 (1992).
 - [6] See, for example, M. Kolář and M. K. Ali, Phys. Rev. A **42**, 7112 (1990).
 - [7] M. Kolář and M. K. Ali, Phys. Rev. A**39**, 6538 (1989).
 - [8] M. Torikai, K. Niizeki, and T. Odagaki, J. Phys. Soc. Jpn **70**, 2917 (2001).
 - [9] J. M. Luck, C. Godrèche, A. Janner, and T. Janssen, J. Phys. A**26**, 1951 (1993).
 - [10] M. Kolář, Phys. Rev. B **47**, 5489 (1993).
 - [11] A. G. Roberts and M. Baake, J. Stat. Phys. **74**, 829 (1994).
 - [12] Z. Y. Wen, F. Wijnands and J. S. W. Lamb, J. Phys. A **27**, 3689 (1994).
 - [13] A. Hopf, O. Knill, and B. Simon, Commun. Math. Phys. **174**, 149(1995).
 - [14] J. Bellissard, Bovier A, and J. -M. Ghez, Rev. Math. Phys. **4**, 1 (1992).
 - [15] However, $\alpha = 1/2$ for the case where E is located at a band edge.
 - [16] J. -P. Allouche and J. Peyrière, C. R. Acad. Sci. Paris **302II**, 1135 (1986).
 - [17] K. Niizeki and R. Endou, to be published.
 - [18] K. Niizeki and E. Yagi, J. Phys. Soc. Jpn **66**, 1387 (1997).
 - [19] A marginal critical state is a peculiar critical state discovered for the case of a ternary SSL in N. Fujita and K. Niizeki, Phys. Rev. Lett. **85**, 4924 (2000). It exhibits a non-power-law scaling.
 - [20] E. L. Albuquerque and M. G. Cottam, Phys. Rep. **376**, 225 (2003).